

## Worksheet 5a: Physics 210: Additional Motion problems

(1) A helicopter is seen to move upward with a position given by:

$$y = y_0 + 4t - 3t^2$$

- (a) What is the velocity of the helicopter at all times?
- (b) What is the vector acceleration of the helicopter at all times?
- (c) If  $y_0=0$ , at what times is the position equal to zero?
- (d) At what time is the velocity zero?

(2) An airplane makes an emergency medical drop at the South Pole. The plane is moving with a velocity of 70 m/s at a distance of 40m above the South Pole.

- (a) How long (in time) before the plane is above the South pole must the package be released in order to hit the South Pole exactly?
- (b) Where was the plane at this time?
- (c) What was the impact velocity vector of the package?

(3) A ball is thrown upward at an angle of  $60^\circ$  with an initial velocity of 30 m/s.

- (a) How long is the ball in flight?
- (b) How high does the ball go above the ground?
- (c) How far in the x-direction does the ball travel?
- (d) What is the impact velocity vector of the ball?

(4) Assume the speed of sound is 343 m/s. A hiker dislodges a rock on the side of a cliff and immediately screams to a hiker below to watch out. If the hiker below is 50 m below the rock, how long will it take between the time that the hiker hears the warning and the rock strikes the position (previous) of the hiker below?

(5) Let's find out how much time is saved at two speeds. Suppose you are going on a 100 mile trip (160934 m). Compare the time that it takes for the trip when driving at two speeds,  $v_1=55$  mph and  $v_2=60$  mph.

While we are talking about driving, what is twice as exciting as 25 m/s in terms of stopping distances? Assume the acceleration in each case is  $-7.62 \text{ m/s}^2$ .

(1) A helicopter is seen to move upward with a position given by:

$$y = y_0 + 4t - 3t^2$$

(a) What is the velocity of the helicopter at all times?

By comparison to the standard free-fall equation we have:

$$v_0 = 4\text{m/s} : -\frac{1}{2}a_y t^2 = -3t^2 \Rightarrow a_y = 6\text{m/s}^2$$

so

$$v = 4 - 6t$$

(b) What is the vector acceleration of the helicopter at all times?

$$\vec{a} = 0\hat{x} - 6\hat{y} \frac{\text{m}}{\text{s}^2}$$

(c) If  $y_0=0$ , at what times is the position equal to zero?

$$0 = 0 + v_{0,y}t - 3t^2 \Rightarrow 0 = t(v_{0,y} - 3t)$$

$$\Rightarrow t = 0 \text{ or } t = \frac{v_{0,y}}{3}$$

(d) At what time is the velocity zero?

$$v_y = v_{0,y} - 6t \Rightarrow t = \frac{v_{0,y}}{6}$$

(2) An airplane makes an emergency medical drop at the South Pole. The plane is moving with a velocity of 70 m/s at a distance of 40m above the South Pole.

(a) How long (in time) before the plane is above the South Pole must the package be released in order to hit the South Pole exactly?

$$y = y_0 + v_{0,y}t - \frac{1}{2}gt^2 \Rightarrow t = \pm\sqrt{\frac{2y}{g}} = 2.86\text{s}$$

(b) Where was the plane at this time?

$$x = v_{0,x}t = 70 \times 2.86 = 200\text{m}$$

(the plane was 200 m from the south pole)

(c) What was the impact velocity vector of the package?

$$v_y = v_{y,0} - gt = -28.02\text{m/s}$$

$$\text{So } \vec{v} = 70\hat{x} - 28\hat{y} \text{ m/s}$$

(3) A ball is thrown upward at an angle of  $60^\circ$  with an initial velocity of 30 m/s.

(a) How long is the ball in flight?

$$v_y = v_{y0} - gt \Rightarrow t = \frac{2v_{y0}}{g} = \frac{2 \times 30 \times \sin(60)}{g} = 5.3\text{s}$$

(b) How high does the ball go above the ground?

$$v_y^2 = v_{y0}^2 - 2g\Delta y \Rightarrow \Delta y = \frac{0^2 - [30\sin(60)]^2}{-2g} = 34.4\text{m}$$

(c) How far in the x-direction does the ball travel?

$$x = x_0 + v_{0,x}t \Rightarrow \Delta x = v_0 \cos(60)t = 79.5\text{m}$$

(d) What is the impact velocity vector of the ball?

$$\vec{v}_f = v_{0,x}\hat{x} - v_{0,y}\hat{y} = 30\cos(60)\hat{x} - 30\sin(60)\hat{y}$$

$$\Rightarrow \vec{v}_f = 15\hat{x} - 26\hat{y} \frac{\text{m}}{\text{s}}$$

(4) Assume the speed of sound is 343 m/s. A hiker dislodges a rock on the side of a cliff and immediately screams to a hiker below to watch out. If the hiker below is 50 m below the rock, how long will it take between the time that the hiker hears the warning and the rock strikes the position (previous) of the hiker below?

$$50 = v_s t_s \Rightarrow t_s = \frac{50}{343} = 0.14s$$

$$-50 = -\frac{1}{2}gt_r^2 \Rightarrow t_r = 3.19s$$

$$T_{\text{to react}} = 3.19s - .14s = 3.05s$$

(5) Let's find out how much time is saved at two speeds. Suppose you are going on a 100 mile trip (160934 m). Compare the time that it takes for the trip when driving at two speeds,  $v_1=55$  mph and  $v_2=60$  mph.

The equation of motion is given for the two speeds. In both cases, the distance traveled is the same.

$$x = v_1 t_1 : x = v_2 t_2$$

Solve these for the times:

$$t_1 = \frac{x}{v_1} : t_2 = \frac{x}{v_2}$$

The difference in times is then:

$$\Delta t = t_2 - t_1 = x \left[ \frac{1}{v_2} - \frac{1}{v_1} \right]$$

As an example, let  $v_1$  be 55 mph (24.6 m/s) and  $v_2$  is 70 mph (31.3 m/s). Then:

$$\Delta t = 100 \left( \frac{1}{70} - \frac{1}{55} \right) = 0.39 \text{hr} = -23 \text{min}$$

How about the same trip at 55 mph and 60 mph?

$$\Delta t = 0.15 \text{hr} = 9 \text{min.}$$

While we are talking about driving, what is twice as exciting as 25 m/s in terms of stopping distances? Assume the acceleration in each case is  $-7.62 \text{ m/s}^2$ .

In terms of stopping distances, assume the accelerations are the same for both cars, but one is at 25 m/s and the other is traveling at such a speed so that the stopping distance is twice that of the first car.

The equations of motion are:

$$0 = v_1^2 - 2a\Delta x_1 : 0 = v_2^2 - 2a\Delta x_2$$

$$\Delta x_2 = 2\Delta x_1$$

$$\Rightarrow v_1^2 - 2a\Delta x_1 = v_2^2 - 2[2a\Delta x_1]$$

$$\Rightarrow v_1^2 - v_2^2 = -2a\Delta x_1$$

So

$$v_2 = \sqrt{v_1^2 + 2a\Delta x_1}$$

This gives the speed at which stopping distances are doubled. To give a more precise answer, let's use some typical values:

Assume the average deceleration rate is  $23 \text{ ft/s}^2$  ( $7.62 \text{ m/s}^2$ ). The car is initially traveling at 25 m/s so the stopping distance is given by:

$$\Delta x_1 = \frac{(25 \text{ m/s})^2}{2(7.62 \text{ m/s}^2)} = 41 \text{m}$$

The speed at which the stopping distance doubles is then:

$$v_2 = \sqrt{2 \times 7.62 \times 2 \times 41} = \sqrt{2}v_1 = 35.4 \text{m/s (about 80 mph).}$$

So twice as exciting as 55 is not 110; it's 80 instead.